

Interstellar H₂ toward HD 37903

Gnaciński Piotr

Institute of Theoretical Physics and Astrophysics, University of Gdańsk,
ul. Wita Stwosza 57, 80-952 Gdańsk
email: pg@iftia9.univ.gda.pl

March 15, 2011

Abstract

We present an analysis of interstellar H₂ toward HD 37903, which is a hot, B 1.5 V star located in the NGC 2023 reflection nebula. Meyer *et al.* (2010) have used a rich spectrum of vibrationally excited H₂ observed by the HST to calculate a model of the interstellar cloud toward HD 37903. We extend Mayer's analysis by including the $\nu''=0$ vibrational level observed by the FUSE satellite.

The T_{01} temperature should not be interpreted as a rotational temperature, but rather as a temperature of thermal equilibrium between the ortho and para H₂. The ortho to para H₂ ratio is lower for the lowest rotational levels than for the higher levels populated by fluorescence. The PDR model of the cloud located in front of HD 37903 points to a gas temperature $T_{kin}=110\text{--}377$ K, hydrogen density $n_H=1874\text{--}544$ cm⁻³ and the star-cloud distance of 0.45 pc.

Key words: *ISM: clouds — ISM: molecules — ultraviolet: ISM*

1 Introduction

A rich spectrum of vibrationally excited H₂ in the direction to HD 37903 was first described by Meyer *et al.* (2001). They have observed over 500 interstellar H₂ absorption lines from excited vibrational levels $\nu''=1\text{--}14$ and rotational levels up to $J''=13$. These lines were detected in a *Hubble Space Telescope* (HST) spectrum made with the *Space Telescope Imaging Spectrograph* (STIS). A *Far Ultraviolet Spectroscopic Explorer* (FUSE) spectrum was made after Mayer's publication allowing to access the $\nu''=0$ vibrational level of the ground electronic state. The FUSE spectrum was used by Rachford *et al.* (2009) to determine the $T_{OP} = 170$ K / $\ln(9N_0/N_1)=68 \pm 7$ K gas "kinetic" temperature and the hydrogen molecular fraction $f(H_2)=2N(H_2)/(2N(H_2)+N(HI))=0.53 \pm 0.09$ in the direction towards HD 37903.

The star HD 37903 was also observed by the *Berkeley*

Extreme and Far-Ultraviolet Spectrometer (BEFS) on-board the *Orbiting and Retrievable Far and Extreme Ultraviolet Spectrometer* (ORFEUS) telescope. The spectral resolution was $R=3000$. Lee *et al.* (2002) have used this spectra to obtain H₂ column densities on $\nu''=0$ and $J''=0\text{--}5$ rotational levels. Their column densities agree within the order of magnitude to the column densities derived in this paper from the FUSE spectra. The physical parameters derived by Lee *et al.* (2002) are: $T_{OP}=63 \pm 5$ K, $f(H_2)=0.496 \pm 0.017$ and the cloud density $n=5600$ cm⁻³.

The H₂ molecule exists in two forms: ortho (odd J'') and para (even J'') H₂. It is caused by the spins of the hydrogen nuclei which can point "in the same" direction (ortho H₂ – triplet state, total nuclei spin $I=1$) or in opposite directions (para H₂ – singlet state, $I=0$). The ratio ortho/para H₂ is therefore 3:1 at standard temperature and pressure. Conversion between this two spin isomers can take place in gas phase, or on the surface of gas grains (Le Boulton 2000). The ortho-para conversion in the gas phase is caused by the exchange of proton in collisions with H, H⁺ and H₃⁺.

The fluorescence cascade of H₂ that leads to population of excited ro-vibrational states of H₂ as well as to the emission of infrared photons from quadrupole transitions has been described by Black & Dalgarno (1976). The first ultraviolet detection of vibrationally excited interstellar H₂ was performed by Federman *et al.* (1995) in the HST spectrum of ζ Ophiuchi.

2 Column densities

We have used both HST STIS and FUSE spectrum to obtain column densities on H₂ ro-vibrational levels. The HST STIS spectrum o59s04010 was averaged because it consist of two subexposures. The HST spectrum is located between 1160 and 1357 Å. We have also used the FUSE observation P1160601. We have analyzed only spectrum from detectors 1B LiF and 1A LiF. These two spectra had the best quality. The FUSE spectra originat-




Figure 1: A fragment of FUSE spectrum with a fit (green line) of H_2 absorption lines. The red line represents continuum.




Figure 2: Fragment of HST STIS spectrum (gray line with dots) fitted with 268 H_2 absorption lines (green line). The figure presents only a 5 Å fragment of the spectrum, while the presented fit was done to the whole 200 Å long STIS spectrum (1160–1357 Å) and included 7449 H_2 lines.




Figure 3: Occupation of H₂ X ro-vibrational levels towards HD 37903. Note the different y-axis scale for $v''=0$ (left) and for higher v'' levels (right).

ing from the same detector were shifted and coadded using the IRAF tasks *pooffsets* and *specalign*. A part of the FUSE spectrum is shown on fig. 1. The whole FUSE spectrum ranges from 987 Å to 1188 Å.

The Ar I 1048 Å line that lies in the wing of R(0) and R(1) lines was cut out from the spectrum. The wings of H₂ lines were calculated up to 60 Å from the line center. The absorption lines in the FUSE spectrum were modeled with the Voigt function, while a Gauss function was used for modeling the H₂ lines in the STIS spectrum.

We have used a Gaussian point spread function (PSF) with FWHM equal to 15 km/s (Jensen *et al.* 2010) for modeling the FUSE spectra. The column densities at $J''=0, 1$ and 2 ($v''=0$) were derived from transitions between B and X electronic levels. The vibrational transitions used are (0,0), (2,0), (3,0) and (4,0), where the first digit is the v' on the upper electronic level B and the second digit $v'' = 0$ is the vibrational level of the ground electronic state X. The FUSE spectrum at the (1,0) vibrational transition was too noisy to perform a good fit. The H₂ line positions and oscillator strengths were adopted from Abgrall *et al.* (1994). The total transition probabilities that include the transitions to continuum (dissociating) states were taken from Abgrall *et al.* (2000).

The column densities of rotational levels $J''=0-9$ from the vibrational $v''=0$ level as well as the column density of the $v''=1$ $J''=0$ level were derived from the FUSE spectrum. Other ro-vibrational levels of the X ground electronic state were derived from the o59s04010 HST STIS spectrum.

The column densities were derived with the profile fitting technique. At each point of the simulated spectrum the optical depth of many spectral lines has been summed. Such procedure was necessary to calculate the profile of blended lines which were ubiquitous in the spectra. The cloud velocity, the doppler broadening parameter and column densities on all observed levels were free parameters, that were fitted to the observed spectra.

The STIS spectrum was fitted with H₂ absorption lines from all vibrational $v''=0-14$ levels, and rotational levels $J''=0-13$. H₂ lines that were blended with atomic lines were excluded from the fitting procedure. Total 7449 H₂ lines were included in the simulated STIS spectrum. The whole STIS spectrum (200 Å long) was fitted at once with all 7449 H₂ lines, because of large number of blended lines. A fragment of the HST STIS spectrum with lines of vibrationally excited molecular hydrogen is presented on Fig. 2. The observed column densities are presented in Table 1 and on Fig. 3. The errors of the column density are about 20% for $J''=0-7$ rotational levels, and up to ~40% for higher rotational levels.




Figure 4: Theoretical $N(J=1)/N(J=0)$ ratio on the $v''=0$ vibrational level from eq. 6 is shown as a dashed blue line. The light blue dashed lines represent errors introduced by the uncertainty of the N_{ortho}/N_{para} H₂ ratio 0.64 ± 0.21 . The observed $N(1)/N(0)$ ratio together with its uncertainty is plotted as a straight line.

3 Results

Total observed column density of H₂ on all vibrational and rotational levels equals to $N(H_2) = (9.6 \pm 1.6) \cdot 10^{20} \text{ cm}^{-2}$. The observed H₂ column density is slightly higher than $N(H_2) = 4 \cdot 10^{20} \text{ cm}^{-2}$ predicted by Meyer *et al.* (2001). Column density of neutral hydrogen $N(HI) = 1.48 \cdot 10^{21} \text{ cm}^{-2}$ (Diplas & Savage 1994). The hydrogen molecular fraction (assuming 10% error of $N(HI)$) equals to $f(H_2) = 2N(H_2)/(2N(H_2) + N(HI)) = 0.56 \pm 0.04$.

The T_{OP} temperature, calculated from the lowest energy levels of ortho and para H₂ ($v'' = 0$) was derived from the equation:

$$\frac{N(1)}{N(0)} = \frac{g_I(1)g_J(1)}{g_I(0)g_J(0)} \exp\left(-\frac{E(1) - E(0)}{kT_{OP}}\right), \quad (1)$$

where g_I is the spin degeneracy factor $g_I(I) = 2I + 1$. The $g_I(1) = 3$ is the statistical weight for ortho H₂, and $g_I(0) = 1$ for the para-H₂ spin isomer. The $g_J(J) = 2J + 1$ is the statistical weight for the rotational level J . The T_{OP} is the temperature of thermal equilibrium between the ortho and para spin isomers. The derived $T_{OP} = 67 \pm 8 \text{ K}$ is similar to the temperatures determined by Rachford *et al.* (2009) (68 K) and by Lee *et al.* (2002) (63 K). The photon dominated regions (PDR) models of interstellar clouds shows that the T_{OP} temperature is correlated with the gas kinetic temperature (Le Petit *et al.* 2006).

However, if we want to calculate the rotational temperature across the ortho-para divide we have to take into account the N_{ortho}/N_{para} H₂ ratio. The population of the ro-vibrational levels of H₂ depends not only from the tem-

Table 1: Column densities of the H₂ ro-vibrational levels towards HD 37903 [cm⁻²].

J''\v''	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
0	5.3e20	7.7e13	2.7e13	1.9e13	1.4e13	1.2e13	6.7e12	8.3e12	4.2e12	3.2e12	2.6e12	2.7e12	1.9e12	1.5e12	1.8e12
1	3.7e20	2.0e14	8.4e13	5.2e13	4.4e13	3.4e13	2.4e13	1.9e13	1.4e13	8.6e12	8.4e12	7.4e12	5.7e12	4.1e12	4.4e12
2	5.6e19	2.3e14	7.4e13	4.6e13	3.8e13	2.7e13	2.7e13	1.5e13	1.2e13	8.6e12	4.2e12	5.3e12	5.3e12	3.7e12	3.8e12
3	6.3e18	2.3e14	8.1e13	5.7e13	3.9e13	3.2e13	2.2e13	1.5e13	1.3e13	8.3e12	6.3e12	5.9e12	5.1e12	3.8e12	2.8e12
4	6.7e17	1.8e14	6.1e13	3.1e13	2.0e13	1.6e13	1.2e13	6.1e12	5.7e12	3.4e12	1.8e12	5.1e12	2.3e12	2.0e12	1.6e12
5	2.6e17	1.1e14	5.3e13	2.9e13	1.9e13	1.2e13	9.7e12	6.0e12	4.2e12	2.7e12	3.5e12	1.8e12	2.1e12	1.4e12	—
6	2.5e16	6.4e13	2.3e13	1.3e13	7.7e12	4.6e12	4.3e12	2.6e12	2.2e12	2.1e12	2.2e12	5.5e11	8.3e11	1.1e12	—
7	8.0e15	5.5e13	2.2e13	1.2e13	8.2e12	7.0e12	3.8e12	3.3e12	1.9e12	2.6e12	1.7e12	1.3e12	1.5e12	8.3e11	—
8	1.9e14	1.3e13	4.4e12	4.0e12	4.3e12	2.2e12	3.1e12	1.8e12	6.0e11	1.4e12	4.6e11	5.0e11	8.4e11	—	—
9	1.6e14	2.7e13	1.0e13	6.4e12	5.9e12	2.7e12	2.4e12	1.2e12	1.0e12	1.4e12	6.8e11	2.0e12	4.3e11	—	—
10	2.3e13	1.1e13	5.0e12	3.0e12	2.0e12	7.8e11	2.2e12	6.2e11	5.8e11	5.6e11	3.2e11	4.5e11	6.7e11	—	—
11	3.3e13	1.7e13	6.2e12	3.6e12	3.2e12	2.9e12	8.5e11	8.1e11	1.1e12	8.9e11	7.1e11	9.5e11	—	—	—
12	4.1e13	7.3e12	1.5e12	4.4e12	1.2e12	3.1e11	8.4e11	1.4e12	6.2e11	3.3e11	2.1e11	2.0e11	—	—	—
13	1.6e13	1.7e13	4.2e12	2.8e12	2.1e12	2.0e12	1.8e12	3.0e11	8.1e11	3.9e11	8.5e11	—	—	—	—



 Figure 5: Rotational temperatures for para and ortho H₂ toward HD 37903 as a function of the vibrational number v'' .

 Table 2: Rotational temperatures for para and ortho H₂ toward HD 37903.

v''	T_{para} [K]	T_{ortho} [K]
1	1962 ± 279	2733 ± 434
2	1779 ± 210	2380 ± 288
3	2086 ± 420	2262 ± 314
4	1701 ± 212	2180 ± 313
5	1337 ± 133	2103 ± 378
6	1613 ± 216	1894 ± 379
7	1536 ± 354	1458 ± 165
8	1318 ± 255	1642 ± 338
9	1194 ± 149	1449 ± 187
10	1009 ± 149	1370 ± 278
11	800 ± 139	1079 ± 278
12	640 ± 127	519 ± 53
13	294 ± 35	305 ± 38
14	92 ± 3	86 ± 9

perature and radiation field, but also from the total ortho to para ratio. In order to include the ortho to para ratio into the Boltzmann equation we will write separate Boltzmann distributions for the ortho and para spin isomers. The Boltzmann distribution for the ortho H₂ can be written as:

$$N_o(J_o'') = \frac{N_{ortho}}{Z_o(T)} g_I(1) g_J(J_o'') \exp\left(-\frac{E(J_o'')}{kT}\right) \quad (2)$$

where N_{ortho} is the total amount of ortho H₂. The partition function Z is:

$$Z_o(T) = \sum_{J_o'' \text{ odd}} g_I(1) g_J(J_o'') \exp\left(-\frac{E(J_o'')}{kT}\right). \quad (3)$$

Let us assume, that para H₂ has a Boltzmann distribution with the same rotational temperature:

$$N_p(J_p'') = \frac{N_{para}}{Z_p(T)} g_I(0) g_J(J_p'') \exp\left(-\frac{E(J_p'')}{kT}\right) \quad (4)$$

where N_{para} is the total amount of para H₂, and the partition function for para H₂:

$$Z_p(T) = \sum_{J_p'' \text{ even}} g_I(0) g_J(J_p'') \exp\left(-\frac{E(J_p'')}{kT}\right). \quad (5)$$

By dividing the equations 2 by 4 we obtain:

$$\frac{N_o(J_o'')}{N_p(J_p'')} = \frac{N_{ortho}}{N_{para}} \frac{g_I(1) g_J(J_o'')}{g_I(0) g_J(J_p'')} \frac{Z_p(T)}{Z_o(T)} \cdot \exp\left(-\frac{E(J_o'') - E(J_p'')}{kT}\right). \quad (6)$$

We have tried to solve the above equation numerically for the $J_o''=1$ and $J_p''=0$ states ($v''=0$) to obtain the T_{01} rotational temperature. The observed $N_{ortho}/N_{para} = 0.64 \pm 0.21$. Figure 4 presents the right side of eq. 6 and the ratio of observed column densities (left side of eq. 6) together with maximal errors. The rotational temperature T_{01} can take values between 0 and 500 K.

The equation 6 can be written in another way by making two assumptions that are **not valid** in the interstellar medium:

1. $T > 240 \text{ K} \Rightarrow Z_p(T)/Z_o(T) \approx 1/3$
2. $N_{ortho}/N_{para} = 3$ as in standard laboratory conditions.

Under this assumptions the eq. 6 for rotational temperature reduces to:

$$\frac{N_o(J_o'')}{N_p(J_p'')} = \frac{g_I(1)g_J(J_o'')}{g_I(0)g_J(J_p'')} \exp\left(-\frac{E(J_o'') - E(J_p'')}{kT}\right). \quad (7)$$

and T_{01} is the same as the temperature T_{OP} of ortho–para thermal equilibrium.

The rotational temperatures T_{02} and T_{13} involve rotational levels of the same spin isomers of H_2 , and do not depend on the total N_{ortho}/N_{para} H_2 ratio and on the partition function. For the cloud towards HD 37903 the $T_{02} = 132 \pm 14 \text{ K}$ and $T_{13} = 172 \pm 9 \text{ K}$. However, the T_{12} temperature calculated from eq. 6 is $133^{+37}_{-25} \text{ K}$, while the use of eq. 7 gives a incorrect temperature of $263 \pm 89 \text{ K}$.

A good illustration that the eq. 6 must be used can be the $\nu=1$ level. On the $\nu = 1$ level the temperature $T_{12} = 1223 \text{ K}$ (levels ortho and para) is close to the $T_{02} = 921 \text{ K}$ and $T_{13} = 1090 \text{ K}$ temperatures calculated from the same spin isomers. The temperature calculated with the traditionally used equation 7 is negative $T = -466 \text{ K}$!

The ortho (odd J'') to para (even J'') H_2 ratio (hereafter O/P) $(O/P)_{\nu''=1-14} = 1.35 \pm 0.18$ was calculated by summing ortho and para H_2 on vibrational levels $\nu''=1-14$. It is a bit lower than $O/P=1.45 \pm 0.08$ given by Meyer *et al.* (2001). We have also used the method proposed by Wilgenbus *et al.* (2000) to calculate O/P H_2 from individual ortho levels, specially $J_o''=1$ and 3 on $\nu''=0$. For the $\nu''=0$ $J''=1$ ortho state the O/P ratio differs significantly from the $(O/P)_{\nu''=1-14} = 1.35$ value and is equal to $(O/P)_{J''=1} = 0.63 \pm 0.11$. Also for the $\nu''=0$ $J''=3$ ortho state the observed O/P is low - only $(O/P)_{J''=3} = 0.70 \pm 0.07$. For higher rotational and vibrational levels the Wilgenbus *et al.* (2000) method gives $(O/P)_{J_o''}$ ratio close to 1.3 derived for excited vibrational H_2 states.

The rotational temperatures for vibrational levels $\nu''=1-14$ for para H_2 were obtained from the linearized Boltzmann distribution:

$$\ln \frac{N_p(J_p'')}{g_I(0)g_J(J_p'')} = \ln N_p(0) - \frac{E(J_p'')}{kT_\nu}. \quad (8)$$

Analogous equation was used for the distribution of the ortho H_2 states. Fig. 3 shows left side of eq. 8 versus E/k . Levels fulfilling the Boltzmann distribution should

Table 3: Comparison of cloud models toward HD 37903.

	this paper	Meyer <i>et al.</i> (2001)	Lee <i>et al.</i> (2002)
telescope	HST STIS and FUSE	HST STIS	BEFS ORFEUS
H_2 levels	$\nu''=0-14$ $J''=0-13$	$\nu''=1-14$ $J''=0-13$	$\nu''=0$ $J''=0-5$
R_V	5.5	5.5	4.1
n_H [cm^{-3}]	544–1874	130–8 800	5 600
d [pc]	0.45	0.5	0.2
T_{kin} [K]	110–377	400	—

be placed on a straight line on this plot. On the $\nu''=0$ vibrational level, which is populated partially by collisions the occupation of rotational levels does not follow a straight line. Therefore the rotational temperature was not calculated for the $\nu''=0$ vibrational level. For both spin isomers and for vibrational levels $\nu''=1-14$ the temperature was calculated using the the linear regression method. The inverse of the line inclination ($-1/T_\nu$ in eq. 8) gives us the rotational temperature. The resulting temperatures are presented in table 2 and on figure 5. The rotational temperatures for the ortho isomer are usually higher than for the para H_2 . However the ortho H_2 rotational temperatures calculated from levels $J_o''=1$ to last but one J_o'' ortho state are in conformity with the para H_2 rotational temperatures. So the assumption that the rotational temperatures T_ν are the same for ortho and para H_2 seems to be justified. The rotational temperatures falls linearly with the increasing vibrational level.

4 Model

We have used the Meudon PDR code (Le Petit *et al.* 2006) to calculate a model of the interstellar medium in the direction of HD 37903. The interstellar reddening $E(B-V)=0.35$ and $R_V=5.5$ was chosen to match the values used by Meyer *et al.* (2001). The radiation source is the HD 37903 star (B 1.5 V spectral type). The model includes 300 H_2 ro-vibrational levels and uses exact computation of radiative transfer in H_2 spectral lines. The model is isobaric with thermal balance.

All our models were two-side models with interstellar radiation field on the observer side equal to one ‘Draine’ unit. The interstellar radiation field on the star side has been varied from 1 to 10^4 ‘Draine’ units. The best model was chosen among 6917 different models by minimising the sum:

$$\sum_{\nu'', J''} w_{\nu'', J''} \left(\log \frac{N_{obs}(\nu'', J'')}{N_{obs}(0, 0)} - \log \frac{N_{model}(\nu'', J'')}{N_{model}(0, 0)} \right)^2 \quad (9)$$

where the weights $w_{\nu'', J''}$ were chosen so, that the levels populated by collisions $J''=0-6$ ($\nu''=0$) have the same in-

fluence on the final sum as the rest of the levels ($v'' > 0$ and $v'' = 0$ $J'' = 7-13$), that are populated by fluorescence.

We have varied the star–cloud distance, hydrogen density and the interstellar radiation field on the star side in order to find a model that matches the observations.

Table 3 presents of comparison of our best model and models presented by Meyer *et al.* (2001) and Lee *et al.* (2002). The star – cloud distance of 0.45 pc, that is responsible for the filling the fluorescence H_2 levels is similar in our model and in the model presented by Meyer *et al.* (2001) ($d=0.5$ pc). Our cloud kinetic temperature (connected with the collisional–filled levels) equals to $T_{kin}=377$ K on the star side of the cloud and $T_{kin}=110$ K on the observer’s side. The hydrogen density obtained in our best model changes from $n_H=544$ cm^{-3} on the star side to 1874 cm^{-3} on the observer’s side. The interstellar radiation field on both sides of our best cloud model is one ‘Draine’ unit.

5 Conclusions

Column density of molecular hydrogen toward HD 37903 on all observed H_2 levels $N(H_2)=9.6 \cdot 10^{20}$ cm^{-2} . The hydrogen molecular fraction $f(H_2)=0.56$. The ortho/para H_2 ratio equals to 1.35 for the excited levels, but for the $J''=1$ ortho state ($v''=0$) the ortho to para ratio is only $(O/P)_{J=1}=0.63$.

The T_{OP} is a temperature of thermal equilibrium between the ortho and para spin isomers and is equal to 67 K. The rotational temperatures $T_{02}=132$ K and $T_{13}=172$ K. The formula for rotational temperatures calculated across the ortho–para divide (like T_{12}) should include the $(O/P)_v$ H_2 ratio.

The best PDR model for the cloud toward HD 37903 gives a gas kinetic temperature $T_{kin}=110-377$ K, hydrogen density $n_H=544-1874$ cm^{-3} and star – cloud distance of 0.45 pc.

Acknowledgments

I would like to thank Herve Abgrall for providing the natural line widths for the H_2 lines. The research was supported by University of Gdańsk grant BW/5400-5-0336-0.

References

Abgrall H., Roueff E., Drira I., 2000, A&ASS, **141**, 297
 Abgrall H., Roueff E., Launay F., Roncin J.-Y., 1994, Can. J. Phys., **72**, 856

Black J.H., Dalgarno A., 1976, ApJ, **203**, 132
 Diplas A., Savage B.D., 1994, ApJSS, **93**, 211
 Draine B.T., Bertoldi F., 1996, ApJ, **468**, 269
 Federman S.R. *et al.*, 1995, ApJ, **445**, 325
 Jensen A., Snow T., Sonneborn G., Rachford B., 2010, ApJ, **711**, 1236
 Le Bourlot J., 2000, A&A, **360**, 656
 Le Petit F., Nehmé C., Le Bourlot J., Roueff E., 2006, ApJSS, **164**, 506
 Lee D.-H. *et al.*, 2002, ApJ, **575**, 234
 Meyer D.M. *et al.*, 2001, ApJ, **553**, L59
 Rachford B.L. *et al.*, 2009, ApJSS, **180**, 125
 Wilgenbus D. *et al.*, 2000, A&A, **356**, 1010